Author Index

Abbott, F.S., see Chan, G.L-Y. 349
An, D.K., see Zhang, X.Y. 343
Ascalone, V.
— and Flamino, L.
Automated high-performance liquid chromatography with column switching for on-line clean-up and analysis of diltiazem and metabolites in human plasma 358
Ashmore, S.P., see Patel, H.R. 269
Au, J.L.-S., see Dalton, J.T. 330
Auriola, S., see Järvinen, T. 13
Averginos, A.
— and Malamataris, S.
High-performance liquid chromatographic determination of indomethacin in human plasma and urine 309
Axelson, J.E., see Chan, G.L.-Y. 349
Bäuerle, A., see Pickert, A. 95
Barchielli, M., see Coppi, G. 245
Barrow, L., see Patel, H.R. 269
Bellotti, V., see Pianezzola, E. 205
Benigni, A., see Chiabrando, C. 1
Bennekom, W.P. van, see Van Opstal, M.A.J. 139
Birder, L.A., see Foglia, J.P. 295
Böswart, J.
—, Schmidt, T., Kostiuk, P., Pacáková, V. and Štulík, K.
High-performance liquid chromatographic determination of some polar phospholipids in serum 61
Botros, H.G.
— and Vijayalakshmi, M.
Cell surface interaction with metal chelates 113
Bruhn, M., see Ellerbroek, L. 314
Bult, A., see Van Opstal, M.AJ. 139
Campoleoni, A., see Chiabrando, C. 1
Castellanos, M., see Wong, B. 21
Chan, G.L.-Y.
—, Axelson, J.E., Abbott, F.S., McErlane, K.M. and Kerr, C.R.
Determination of 5-hydroxypropafenone in biological fluids by fused-silica capillary gas chromatography using electron-capture detection 349
Chan, K.K., see Hong, P.S. 131
Chasseaud, L.F., see Taylor, I.W. 275
Chen, J., see Zhang, X.Y. 343
Chiabrando, C.
—, Pinciroli, V., Campoleoni, A., Begnini, A., Piccinelli, A. and Fanelli, R.
Quantitative profiling of 6-ketoprostaglandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinor-thromboxane B_2 in human and rat urine by immunoaffinity extraction with gas chromatography–mass spectrometry 1
Colussi, R.D., see Sioufi, A. 195
Coppi, G.
— and Barchielli, M.
Simple method for the determination of L-5-hydroxytryptophan in plasma by high-performance liquid chromatography 245
Dalton, J.T.
—, Geuns, E.R. and Au, J.L.-S.
High-performance liquid chromatographic determination of mitomycin C in rat and human plasma and urine 330
Desai, D.M., see Pianezzola, E. 205
Dubois, J.P., see Sioufi, A. 195
Durrer, A.
—, Walther, B., Racciatti, A. and Testa, B.
Convenient method for the analysis of nicotinic acid as a metabolite of nicotinate esters in various tissue homogenates 256
Edwards, G., see Idowu, O.R. 167
Elderfield, A.J.
—, Truscott, R.J.W., Gan, I.E.T. and Schier, G.M.
Separation of tryptophan metabolites by reversed-phase high-performance liquid chromatography with amperometric and fluorescence detection 71
Ellerbroek, L.
— and Bruhn, M.
Determination of flumequine in biological fluids and meat by high-performance liquid chromatography 314
Elliger, S.S.
— and Watson, G.
Method for the determination of the specific activities of UTP and CTP in mouse kidney by high-performance liquid chromatography 249
Fanelli, R., see Chiabrando, C. 1

Ferrario, P., see Zecca, L. 303
Flaminio, L., see Ascalone, V. 358
Foglia, J.P.
—, Birder, L.A. and Perel, J.M.
Determination of fluvoxamine in human plasma by high-performance liquid chromatography with ultraviolet detection 295
Fontana, E., see Pianezzola, E. 205
Gal, J., see Pianezzola, E. 205
Gan, I.E.T., see Elderfield, A.J. 71
Geuns, E.R., see Dalton, J.T. 330
Governa, M., see Marchiseppe, I. 288
Gröhler, G., see Krause, W. 324
Hamoir, T., see Musch, G. 215
Haymond, M.W., see Horber, F.F. 81
Henion, J.D., see Mück, W.M. 41
Herbert, K., see James, I.T. 105
Hikida, K.
—, Inoue, Y., Miyazaki, T., Kojima, N. and Ohkura, Y.
Determination of bromperiodol in serum by automated column-switching high-performance liquid chromatography 227
Hirvonen, E., see Järvinen, T. 13
Holá, V., see Jelínek, I. 338
Holthuis, J.J.M., see Van Opstal, M.A.J. 139
Hong, P.S.
— and Chan, K.K.
Analysis of 4-hydroxycyclophosphamide by gas chromatography–mass spectrometry in plasma 131
Horber, F.F.
—, Kahl, J., Lecavalier, L., Krom, B. and Haymond, M.W.
Determination of leucine and α-ketoisocaproic acid concentrations and specific activity in plasma and leucine specific activities in proteins using high-performance liquid chromatography 81
Horst, F.A.L. van der, see Van Opstal, M.A.J. 139
Idowu, O.R.,
—, Ward, S.A. and Edwards, G.
Determination of artelinic acid in blood plasma by high-performance liquid chromatography 167
Inoue, Y., see Hikida, K. 227
Jacqz-Aigrain, E.
—, Menard, Y., Popon, M. and Mathieu, H.
Dextromethorphan phenotypes determined by high-performance liquid chromatography and fluorescence detection 361
Järvinen, T.
—, Keinonen, T., Auriola, S., Peura, P., Hirvonen, E. and Palva, E.
Specific and sensitive quantitation of medroxyprogesterone acetate in human serum by gas chromatography–mass spectrometry 13
James, I.T.
—, Herbert, K., Perrett, D. and Thompson, P.W.
Determination of serum cytidine deaminase activity using ion-pair reversed-phase liquid chromatography 105
Jarvi, E.J., see Patrick, K.S. 123
Jelínek, I.
—, Rejholec, V., Holá, V., Roubal, Z., Snopek, J. and Smolková-Keulemansová, E.
Isotachophoretic determination of sulbactam in rat serum 338
Kahl, J., see Horber, F.F. 81
Kasashima, K., see Kawai, S. 235
Kawai, S.
—, Kasashima, K. and Tomita, M.
High-performance liquid chromatographic determination of malonaldehyde in serum 235
Keinonen, T., see Järvinen, T. 13
Kerr, C.R., see Chan, G.L.-Y. 349
Koel, M.
— and Nebinger, P.
Simultaneous automated high-performance liquid chromatographic determination of 5-hydroxy-3-indoleacetic acid and homovanillic acid in urine with fluorescence detection 263
Kojima, N., see Hikida, K. 227
Kostiuk, P., see Böswart, J. 61
Krause, W.
— and Gröhler, G.
Automated high-performance liquid chromatographic assay for the β-carboline derivative Abecarnil in plasma 324
Krom, B., see Horber, F.F. 81
LaCreta, F.P., see Tinsley, P.W. 318
Lampert, B.M.
— and Stewart, J.T.
Determination of cocaine and selected metabolites in canine and human serum

by reversed-phase high-performance liquid chromatography on coupled cyanopropyl and silica columns 153
Lecavalier, L., see Horber, F.F. 81
Liebich, H.M., see Picker, A. 95
Lohmar, F.
—, Musch, E., Von Unruh, G.E. and Mühlenbruch, B.J.
Gas chromatographic assay for chlorambucil and phenylacetic mustard in plasma 281
McErlane, K.M., see Chan, G.L.-Y. 349
Makin, H.L.J., see Street, J.M. 31
Malamataris, S., see Avgerinos, A. 309
Marchiseppe, I.
—, Valentino, M., Governa, M. and Stocchi, V.
Determination of total 2,5-hexanedione by reversed-phase high-performance liquid chromatography 288
Marfil, F., see Sioufi, A. 195
Massart, D.L., see Musch, G. 215
Mathieu, H., see Jacqz-Aigrain, E. 361
Matsubayashi, K.
—, Une, T. and Osada, Y.
Determination of ofloxacin in bronchoalveolar lavage fluid by high-performance liquid chromatography and fluorimetric detection 354
Menard, Y., see Jacqz-Aigrain, E. 361
Meyer, M.C., see Patrick, K.S. 123
Miyazaki, T., see Hikida, K. 227
Moro, E., see Pianezzola, E. 205
Mück, W.M.
— and Henion, J.D.
Determination of leucine enkephalin and methionine enkephalin in equine cerebrospinal fluid by microbore high-performance liquid chromatography and capillary zone electrophoresis coupled to tandem mass spectrometry 41
Mühlenbruch, B.J., see Lohmar, F. 281
Musch, E., see Lohmar, F. 281
Musch, G.
—, Hamoir, T. and Massart, D.L.
Determination of pentoxifylline and its 5-hydroxy metabolite in human plasma by solid-phase extraction and high-performance liquid chromatography with ultraviolet detection 215
Nancollas, G.H., see Singh, R.P. 239
Nebinger, P., see Koel, M. 263
O'Dwyer, P.J., see Tinsley, P.W. 318
Ohkura, Y., see Hikidai, K. 227
Opstal, M.A.J. van, see Van Opstal, M.A.J. 139
Osada, Y., see Matsubayashi, K. 354
Pacáková, V., see Böswart, J. 61
Palva, E., see Järvinen, T. 13
Patel, H.R.
—, Ashmore, S.P., Barrow, L. and Tanner, M.S.
Improved high-performance liquid chromatographic technique for the determination of hepatic α-tocopherol 269
Patrick, K.S.
—, Jarvi, E.J., Straughn, A.B. and Meyer, M.C.
Gas chromatographic–mass spectrometric analysis of plasma nifedipine 123
Perel, J.M., see Foglia, J.P. 295
Perrett, D., see James, I.T. 105
Peura, P., see Järvinen, T. 13
Pianezzola, E.
—, Bellotti, V., Fontana, E., Moro, E., Gal, J. and Desai, D.M.
Determination of the enantiomeric composition of salsolinol in biological samples by high-performance liquid chromatography with electrochemical detection 205
Piccinelli, A., see Chiabrando, C. 1
Pickert, A.
—, Bäuerle, A. and Liebich, H.M.
Determination of hippuric acid and furanic acid in serum of dialysis patients and control persons by high-performance liquid chromatography 95
Pinciroli, V., see Chiabrando, C. 1
Popon, M., see Jacqz-Aigrain, E. 361
Racciatti, A., see Durrer, A. 256
Rejholec, V., see Jelínek, I. 338
Richard, J., see Sioufi, A. 195
Roubal, Z., see Jelínek, I. 338
Schier, G.M., see Elderfield, A.J. 71
Schmidt, T., see Böswart, J. 61
Singh, R.P.
—, Smesko, S.A. and Nancollas, G.H.
Simultaneous determination of citrate and D-isocitrate in urine by isocratic ion chromatography 239

Sioufi, A.
—, Marfil, F., Richard, J., Colussi, D. and Dubois, J.P.
High-performance liquid chromatographic determination of pirprofen and five of its metabolites in human plasma without hydrolysis and in human urine before and after chemical hydrolysis 195
Smesko, S.A., see Singh, R.P. 239
Smolková-Keulemansová, E., see Jelínek, I. 338
Snopek, J., see Jelínek, I. 338
Stewart, J.T., see Lampert, B.M. 153
Stocchi, V., see Marchiseppe, I. 288
Straughn, A.B., see Patrick, K.S. 123
Street, J.M.
—, Trafford, D.J.H. and Makin, H.L.J.
Carrier gas flow-rate and injection system in capillary gas chromatography of unconjugated and glycine-conjugated bile acids 31
Streete, P.J.
Rapid high-performance liquid chromatographic methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Štulík, K., see Böswart, J. 61
Tanner, M.S., see Patel, H.R. 269
Taylor, I.W.
— and Chasseaud, L.F.
Determination of ximoprofen in human plasma by gas chromatography 275
Testa, B., see Durrer, A. 256
Thompson, P.W., see James, I.T. 105
Tinsley, P.W.
—, O'Dwyer, P.J. and LaCreta, F.P.
High-performance liquid chromatographic analysis of N,N′,N′-triethylenethiophosphoramide in human plasma 318
Tomita, M., see Kawai, S. 235
Trafford, D.J.H., see Street, J.M. 31
Truscott, R.J.W., see Elderfield, A.J. 71
Une, T., see Matsubayashi, K. 354
Unruh, G.E. Von, see Lohmar, F. 281
Valentino, M., see Marchiseppe, I. 288
Van Bennekom, W.P., see Van Opstal, M.A.J. 139
Van der Horst, F.A.L., see Van Opstal, M.A.J. 139
Van Opstal, M.A.J.
—, Van der Horst, F.A.L., Holthuis, J.J.M., Van Bennekom, W.P. and Bult, A.
Automated reversed-phase chromatographic analysis of etoposide and teniposide in plasma by using on-line surfactant-mediated sample clean-up and column-switching 139
Vijayalakshmi, M., see Botros, H.G. 113
Von Unruh, G.E., see Lohmar, F. 281
Walther, B., see Durrer, A. 256
Ward, S.A., see Idowu, O.R. 167
Watson, G., see Elliger, S.S. 249
Wong, B.
— and Castellanos, M.
Enantioselective measurement of the *Candida* metabolite D-arabinitol in human serum using multi-dimensional gas chromatography and a new chiral phase 21
Wu, R.J., see Zhang, X.Y. 343
Zecca, L.
— and Ferrario, P.
Determination of diclofenac in plasma and synovial fluid by high-performance liquid chromatography with electrochemical detection 303
Zhang, X.Y.
—, Wu, R.J., Chen, J. and An, D.K.
Determination of glycyrrhizin and its metabolite glycyrrhetinic acid in rabbit plasma by high-performance liquid chromatography after oral administration of licrozin 343

Subject Index

Abecarnil
Automated HPLC assay for the β-carboline derivative Abecarnil in plasma 324
Aldophosphamide
Analysis of 4-hydroxycyclophosphamide by GC–MS in plasma 131
Anthranilic acid
Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71
D-Arabinitol
Enantioselective measurement of the *Candida* metabolite D-arabinitol in human serum using multi-dimensional GC and a new chiral phase 21
Artelinic acid
Determination of artelinic acid in blood plasma by HPLC 167
Benzoylecgonine
Determiantion of cocaine and selected metabolites in canine and human serum by reversed-phase HPLC on coupled cyanopropyl and silica columns 153
Benzoylnorecgonine
Determination of cocaine and selected metabolites in canine and human serum by reversed-phase HPLC on coupled cyanopropyl and silica columns 153
Bile acids
Carrier gas flow-rate and injection system in capillary GC of unconjugated and glycine-conjugated bile acids 31
Bromperidol
Determination of bromperidol in serum by automated column-switching HPLC 227
Cell surface interactions
Cell surface interactions with metal chelates 113
Chlorambucil
GC assay for chlorambucil and phenylacetic mustard in plasma 281
Citrate
Simultaneous determination of citrate and D-isocitrate in urine by isocratic ion chromatography 239
Cocaine
Determination of cocaine and selected metabolites in canine and human serum by reversed-phase HPLC on coupled cyanopropyl and silica columns 153
CTP
Method for the determination of the specific activities of UTP and CTP in mouse kidney by HPLC 249
Cytidine
Determination of serum cytidine deaminase activity using ion-pair reversed-phase LC 105
Cytidine deaminase
Determination of serum cytidine deaminase activity using ion-pair reversed-phase LC 105
Deacetyldiltiazem
Automated HPLC with column switching for on-line clean-up and analysis of diltiazem and metabolites in human plasma 358
Deacetyl-N-monodemethyldiltiazem
Automated HPLC with column switching for on-line clean-up and analysis of diltiazem and metabolites in human plasma 358
Dextromethorphan
Dextromethorphan phenotypes determined by HPLC and fluorescence detection 361
Dextrorphan
Dextromethorphan phenotypes determined by HPLC and fluorescence detection 361
Diclofenac
Determination of diclofenac in plasma and synovial fluid by HPLC with electrochemical detection 303
Diflunisal
Rapid HPLC methods for the detemination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Diltiazem
Automated HPLC with column switching for on-line clean-up and analysis of diltiazem and metabolites in human plasma 358
2,3-Dinor-6-ketoprostaglandin $F_{1\alpha}$
Quantitative profiling of 6-ketoprosta-

glandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinor-thromboxane B_2 in human and rat urine by immunoaffinity extraction with GC–MS 1

2,3-Dinor-thromboxane B_2

Quantitative profiling of 6-ketoprostaglandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinor-thromboxane B_2 in human and rat urine by immunoaffinity extraction with GC–MS 1

Enkephalins

Determination of leucine enkephalin and methionine enkephalin in equine cerebrospinal fluid by microbore HPLC and capillary zone electrophoresis coupled to tandem MS 41

Enzymes

Determination of serum cytidine deaminase activity using ion-pair reversed-phase LC 105

Etoposide

Automated reversed-phase chromatographic analysis of etoposide and teniposide in plasma by using on-line surfactant-mediated sample clean-up and column-switching 139

Fenoprofen

Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179

Flumequine

Determination of flumequine in biological fluids and meat by HPLC 314

Fluvoxamine

Determination of fluvoxamine in human plasma by HPLC with UV detection 295

Furanic acid

Determination of hippuric acid and furanic acid in serum of dialysis patients and control persons by HPLC 95

Glycine-conjugated bile acids

Carrier gas flow-rate and injection system in capillary GC of unconjugated and glycine-conjugated bile acids 31

Glycyrrhetinic acid

Determination of glycyrrhizin and its metabolite glycyrrhetinic acid in rabbit plasma by HPLC after oral administration of licorzin 343

Glycyrrhizin

Determination of glycyrrhizin and its metabolite glycyrrhetinic acid in rabbit plasma by HPLC after oral administration of licorzin 343

2,5-Hexanedione

Determination of total 2,5-hexanedione by reversed-phase HPLC 288

Hippuric acid

Determination of hippuric acid and furanic acid in serum of dialysis patients and control persons by HPLC 95

Homovanillic acid

Simultaneous automated HPLC determination of 5-hydroxy-3-indoleacetic acid and homovanillic acid in urine with fluorescence detection 263

Hydroxyanthranilic acids

Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71

4-Hydroxycyclophosphamide

Analysis of 4-hydroxycyclophosphamide by GC–MS in plasma 131

5-Hydroxy-3-indoleacetic acid

Simultaneous automated HPLC determination of 5-hydroxy-3-indoleacetic acid and homovanillic acid in urine with fluorescence detection 263

3-Hydroxykynurenine

Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71

5-Hydroxypentoxifylline

Determination of pentoxifylline and its 5-hydroxy metabolite in human plasma by solid-phase extraction and HPLC with UV detection 215

Hydroxypropafenone

Determination of 5-hydroxypropafenone in biological fluids by fused-silica capillary GC using electron-capture detection 349

5-Hydroxytryptophan

Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71

L-5-Hydroxytryptophan

Simple method for the determination of

L-5-hydroxytryptophan in plasma by HPLC 245
Ibuprofen
Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Indomethacin
Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Indomethacin
HPLC determination of indomethacin in human plasma and urine 309
D-Isocitrate
Simultaneous determination of citrate and D-isocitrate in urine by isocratic ion chromatography 239
α-Ketoisocaproic acid
Determination of leucine and α-ketoisocaproic acid concentrations and specific activity in plasma and leucine specific activities in proteins using HPLC 81
Ketoprofen
Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
6-Ketoprostaglandin $F_{1\alpha}$
Quantitative profiling of 6-ketoprostaglandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinorthromboxane B_2 in human and rat urine by immunoaffinity extraction with GC–MS 1
Kynurenine
Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71
Leucine
Determination of leucine and α-ketoisocaproic acid concentrations and specific activity in plasma and leucine specific activities in proteins using HPLC 81
Leucine enkephalin
Determination of leucine enkephalin and methionine enkephalin in equine cerebrospinal fluid by microbore HPLC and capillary zone electrophoresis coupled to tandem MS 41
Malonaldehyde
HPLC determination of malonaldehyde in serum 235
Medroxyprogesterone acetate
Specific and sensitive quantitation of medroxyprogesterone acetate in human serum by GC–MS 13
Mefenamic acid
Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Metal chelates
Cell surface interactions with metal chelates 113
Methionine enkephalin
Determination of leucine enkephalin and methionine enkephalin in equine cerebrospinal fluid by microbore HPLC and capillary zone electrophoresis coupled to tandem MS 41
Mitomycin C
HPLC determination of mitomycin C in rat and human plasma and urine 330
N-Monodemethyldiltiazem
Automated HPLC with column switching for on-line clean-up and analysis of diltiazem and metabolites in human plasma 358
Naproxen
Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Nicotinate esters
Convenient method for the analysis of nicotine acid as a metabolite of nicotinate esters in various tissue homogenates 256
Nicotinic acid
Convenient method for the analysis of nicotinic acid as a metabolite of nicotinate esters in various tissue homogenates 256
Nifedipine
GC–MS analysis of plasma nifedipine 123
Non-steroidal anti-inflammatory drugs
Rapid HPLC method for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179

Norcocaine
Determination of cocaine and selected metabolites in canine and human serum by reversed-phase HPLC on coupled cyanopropyl and silica columns 153
Nucleotides
Method for the determination of the specific activities of UTP and CTP in mouse kidney by HPLC 249
Ofloxacin
Determination of ofloxacin in bronchoalveolar lavage fluid by HPLC and fluorimetric detection 354
Pentoxifylline
Determination of pentoxifylline and its 5-hydroxy metabolite in human plasma by solid-phase extraction and HPLC with UV detection 215
Phenylacetic mustard
GC assay for chlorambucil and phenylacetic mustard in plasma 281
Phosphatidylcholine
HPLC determination of some polar phospholipids in serum 61
Phosphatidylethanolamine
HPLC determination of some polar phospholipids in serum 61
Phospholipids
HPLC determination of some polar phospholipids in serum 61
Piroxicam
Rapid HPLC methods for the determination of overdose concentrations of some non-steroidal anti-inflammatory drugs in plasma or serum 179
Pirprofen
HPLC determination of pirprofen and five of its metabolites in human plasma without hydrolysis and in human urine before and after chemical hydrolysis 195
Prostaglandins
Quantitative profiling of 6-ketoprostaglandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinorthromboxane B_2 in human and rat urine by immunoaffinity extraction with GC–MS 1
Salsolinol
Determination of the enantiomeric composition of salsolinol in biological samples by HPLC with electrochemical detection 205
Sphingomyelin
HPLC determination of some polar phospholipids in serum 61
Steroids
Specific and sensitive quantitation of medroxyprogesterone acetate in human serum by GC–MS 13
Sulbactam
Isotachophoretic determination of sulbactam in rat serum 388
Teniposide
Automated reversed-phase chromatographic analysis of etoposide and teniposide in plasma by using on-line surfactant-mediated sample clean-up and column-switching 139
Thromboxane B_2
Quantitative profiling of 6-ketoprostaglandin $F_{1\alpha}$, 2,3-dinor-6-ketoprostaglandin $F_{1\alpha}$, thromboxane B_2 and 2,3-dinorthromboxane B_2 in human and rat urine by immunoaffinity extraction with GC–MS 1
α-Tocopherol
Improved HPLC technique for the determination of hepatic α-tocopherol 269
N,N′,N′′-Triethylenethiophosphoramide
HPLC analysis of N,N′,N′′-triethylenethiophosphoramide in human plasma 318
Tryptophan
Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71
Uridine
Determination of serum cytidine deaminase activity using ion-pair reversed-phase LC 105
UTP
Method for the determination of the specific activities of UTP and CTP in mouse kidney by HPLC 249
Vitamin E
Improved HPLC technique for the determination of hepatic α-tocopherol 269
Xanthurenic acid
Separation of tryptophan metabolites by reversed-phase HPLC with amperometric and fluorescence detection 71
Ximoprofen
Determination of ximoprofen in human plasma by GC 275